

BROCCOLI

Harvest Guide 2024 Lockyer Valley Queensland





"Buttoning "or premature flowering can occur in young broccoli plants primarily in the spring timeslot. This is caused by stress to the seedlings either at the nursery stage or soon after transplanting from a range of factors including exposure to cold temperatures, extra/rapid hardening of the plants and wet/dry soil conditions. This problem is more pronounced in districts with cooler late winter/spring temperatures. Some varieties are more susceptible than others. However, even varieties that normally would not present with a significant issue in most years could still show symptoms under some conditions as noted. "Blindness" is a condition whereby the growing point of a seedling/young plant is damaged thereby preventing the production of a marketable head. Several factors can cause this including low and high temperatures, insect damage as well as non-recommended high levels of chemicals or fertilisers. At times, symptoms may not express in the nursery but rather in the field, several weeks after transplanting.

Ross Edser | 0419 709 626 David Commens | 0419 726 826 Luke Murphy-Clarke | 0418 158 079

The information provided is based on an average of data and observations collected from our trials. Significant variations may occur in the performance due to a range of conditions including cultural/management practices, climate, soil type and geographic location. As a consequence South Pacific Seeds cannot accept any liability as to the accuracy of this information. VALID FOR 12 MONTHS FROM NOVEMBER 2023. ACN 002 887 256

Griffith | 02 6962 7333 Adelaide | 0417 545 099 Brisbane | 07 3393 3766 Melbourne | 03 9770 8441 Perth | 08 9331 6356

DIRECT SOWING: Allow an extra 2-3 weeks depending on the time of year.

Harvest

Transplant

